

REVIEW OF USE OF ALTERNATIVES TO METHYL BROMIDE FOR HORTICULTURAL CROPS IN ASIA

J.G.M. Vos* and G.S. Lim

Methyl bromide is a broad spectrum pesticide, principally used as a fumigant. On soil, it is used as a pre-plant soil fumigant in locations where a broad complex of soil-borne pests limit the production of certain crops, particularly when they are repeatedly grown on the same land. In SE Asia and the Pacific, the use of methyl bromide as a soil sterilant is only about 6 % of all methyl bromide used (UNDP, 1995). Alternative methods to control soil pests are widespread. Aware of the negative side effects of pesticides, alternative soil pest control practices with reduced chemical inputs are being suggested. Soil pest control practices include biological control by means of natural enemies and integrated methods, such as integrated pest management (IPM) or integrated crop management (ICM). This paper documents a number of cases of alternative soil pest control methods with reduced chemical inputs in commercial use for horticultural crops in Asia. In addition, alternatives that have been found effective in field and/or greenhouse trials are reviewed.

Cases of alternative soil pest control methods used commercially for horticultural crops on modern as well as traditional farms in Asia are listed in Table 1. The methods range from relatively new techniques, such as substrates and solarisation, to improved methods of traditional techniques, such as composting. There is now greater appreciation and respect for low-input, traditional agricultural systems, and the methods that are employed by traditional farmers are recognised as often being more efficient and logical than previously thought. In fact, to manage nematodes, integrated systems are being developed that incorporate or consider many alternative small-scale farmer practices (Bridge, 1996). Most of the identified techniques can be applied in intensive horticultural production.

In addition to the cases of commercially applied alternative soil pest control methods described in the previous section, there are numerous reports of IPM experimental efforts and successes in many parts of the world, and farmers would benefit from further verification on a larger scale at the farm level. Table 2 lists cases of experimental use of soil pest control methods that could be used as elements of an IPM / ICM approach to successfully replace methyl bromide, with a general indication of applicability.

This review shows that alternative methods to manage soil-borne pests are widespread on various crops in Asia. The choice of method(s) varies to suit the cropping system and the environment. It is concluded that practices do exist to manage soil-borne pests successfully in ecologically sound ways to achieve sustainable crop production.

It should be stressed that, in addition to the documented cases given here, many other (cultural) practices are widespread throughout the Asian region which are applied for various reasons. These, integrated with other crop management measures contribute to the management of soil pests in general, such as sanitation, various methods of tillage (e.g. repeated summer ploughing), crop rotation, balanced fertilising, regulation of irrigation, optimising planting time, choice of seed / planting material, planting distance and habitat management. IPM integrates the available pest control methods to achieve a farmer's most effective, economical, and sustainable combination for a particular local situation. There are now many IPM success cases reported at the farm level for a variety of crops, including horticultural crops. In Asia, all of these success cases contradict the claim that methyl bromide would be needed to grow commercial horticultural crops successfully and economically.

Table 1: Cases using alternative practices for the effective control of soil pests commercially.

Crop(s)	Method	Country/ies	Source(s) of information
1 Various horticultural crops	Composting and using compost	India Malaysia Philippines South Korea Thailand Elsewhere	IIBC staff Harris (1996) IFOAM (1995)
2 Shallot Hot pepper Tomato Other vegetables	Flooding of fields	Indonesia Malaysia Vietnam Elsewhere	IIBC staff Vos et al. (1993) Sariah and Tanaka (1994) Vu (1990)
3 Strawberry Tomato Pepper Leafy vegetables	Cultivation on substrates	Malaysia Indonesia Singapore	IIBC staff Various. private growers
4 Cabbage Solanaceous crops	Application of lime	Malaysia Thailand Philippines Elsewhere	IIBC staff Vattanatangum (1990)
5 Pepper Tomato Watermelon	Application of mulch	Taiwan Indonesia Malaysia Vietnam	IIBC staff Vos and Sumami (1997) VU(1990).
6 Cabbage	Treatment of seed	Countries in Asia	IIBC staff Vattanatangum (1990)
7 Various horticultural crops	Burning of top-soil	India Bangladesh	IIBC staff Choudhury and Hoque (1982)
8 Various horticultural crops	Application of neem cake	India	IIBC staff Alarn (1991)
9 Various horticultural crops	Mixed cropping with marigold	India Vietnam	IIBC staff Khan et al. (1971)
10 Various horticultural crops	Tillage of soil	Vietnam Bangladesh India	IIBC staff Vu(1990) Hossein (1990) Prasad and Chawla (1991)
11 Cucumber Eggplant Tomato	Grafting on resistant root stocks	China Japan	IIBC: staff Qiu (1990)
12 Various horticultural crops	Use of resistant cultivars	Japan Malaysia Elsewhere	IIBC staff Takeuchi (1990)
13 Various horticultural crops	Solarisation in greenhouses	Japan	Horiuchi (1991)
14 Potato	Increase of hilling and irrigation	Australia	Ansari (1990)
15 Tomato	Rotation of crops	Bangladesh	Hossain (1990)
16 Various horticultural crops	Solarisation of seed beds	China	Qiu (1990)
17 Tomato Eggplant Other vegetables	Adjusting of planting time	India	Sethi and Gaur (1986)

Table 2: Alternative soil post control methods found effective in tests or trials.

Crop(s)	Method	Country/ies	Source(s) of information	Applicability
1 Tomato	Resistance against bacterial wilt	Indonesia Elsewhere	Purwati and Hanudin (1995)	Regions with equal wilt strains
2 Cabbage Onion	Solarisation	Philippines Australia	Boteng (1990) Porter and Merriman (1985)	Regions with hot sunny climates
3 Eggplant	Biological control using <i>Talaromyces flavus</i>	USA	Marois et al. (1982)	Fields and greenhouses
4 Cumin Clusterbean	Solarisation, summer irrigation, amendmets	India	Lodha (1995)	Hot add regions
5 Tomalo Strawberry	Integration of biological and chemical control	Israel	Elad et al. (1994)	Fields and greenhouses with <i>Botrytis cinerea</i>
6 Tomato	Resistance to rootknot nematode	Argentina Elsewhere	Gallardo (1988)	Tomato growing areas
7 Lettuce	Biological control using <i>Trichoderma viride</i>	Austria	Bedlan (1988)	Greenhouse lettuce crops
8 Cucumber Tomato	Integration of biological and cultural control	Japan	Kobayashi (1991)	Vegetable crops in greenhouses and fields
9 Tomato	Use of trap crops	India	Rangaswarny and Reddy (1993)	Greenhouses and fields
10 Tomato Cucumber rootstocks Eggplant Pepper	Grafting on resistant	Switzerland Japan China Italy Korea	Dufour and Taillens (1994) Kuwata et al. (1994) Lu eta/. (1992) Morra eta/. (1992) Choe (1989)	Fields and greenhouses
11 Banana	Hot water treatment	?	Prasad and Reddy (1994)	Small-hold farms
12 Tea	Soil sterilisation for young tea plants	India	Rao (1976)	Small-hold farms

References:

- Alam, M.M. (1991)
Control of plant-parasitic nematodes in field Brinjal. *Journal of Nematology* 9: p. 21-30
- Bedlan, G. (1988)
Bekämpfungsmöglichkeiten der Schwarzfaule mit Hilfe des Antagonisten *Trichoderma viride* Pers. an Salat im Freiland. *Pflanzenschutz* 4: p. 8-10
- Boteng, M.D.C. (1990) Control of clubroot (*Plasmodiophora brassicae*) on cabbage (*Brassica oleracea*). B.S. thesis, Benguet State University, La Trinidad, Benguet, Philippines: 27 pp.
- Bridge, J. (1996) Nematode management in sustainable and subsistence agriculture. *Annual Review of Phytopathology* 34: p. 201-225.
- Choe, J.S. (1989)
Phytophthora blight of green pepper in Korea. Extension bulletin ASPAC, Food and fertilizer technology centre 302: p. 18-25
- Choudhury, B.C. and Hoque, M.O. (1982)
Control of root-knot nematode in brinjal seed beds by heat treatments. *Bangladesh Journal of Agricultural Research* 7: p. 55-57
- Dufour, L. and Taillens, J. (1994)
L'effacement des nématodes hémiparasites de la tomate et la pourriture noire des racines. *Revue Suisse de Viticulture* 26(2): p. 85-88
- Elad, Y., Shtienberg, D. and Niv, A. (1994) *Trichoderma harzianum* T39 integrated with fungicides: improved biocontrol of grey mould. Brighton crop protection conference - pests and diseases 1994: p. 1109-1114
- Gallardo, G.S. (1988) Breeding of processing tomatoes with genetic resistance to rootknot nematodes (*Meloidogyne incognita*). Proceedings of the international symposium on integrated management practices in tomato and pepper production in the tropics, Tainan, Taiwan. AVRDC. p. 107-112
- Harris, P.J.C. (1996) An assessment of the potential contribution of organic expertise and techniques to the development of sustainable farming systems for the Eastern India Rainfed Farming Project. Consultancy report, Henry Doubleday Research Association, UK: 45 pp.
- Horiuchi, S. (1991)
Solarisation for greenhouse crops in Japan. In: DeVay, J.E., Stapleton, J.J. and Elmore, C.L. (eds) Soil solarisation. Proceedings of the first international conference on soil solarisation. FAO plant production and protection paper 109: p. 16-27
- Hossain, Q.T. (1990)
Status and management of vegetable pests in Bangladesh. In: Lim, G.S. and Di, Y.-B. (eds) Status and management of major vegetable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 23-29
- IFOAM (1995)
IFOAM conference on food, culture, trade and the environment, 19-22 July 1995, Seoul, Korea.

- Khan, A.M., Saxena, S.K. and Siddiqi, Z.A. (1971) Efficacy of *Tagetes erecta* in reducing root infesting nematodes of tomato and okra. *Indian Phytopathology* 24: p. 166-169.
- Kobayashi, N. (1991) Biological control of soil-borne diseases with VAM fungi and charcoal compost. In: Komada, H, Kiritani, K. and Bay-Petersen, J. (eds) *The biological control of plant diseases. Proceedings of the international seminar Tsukuba, Japan, 17-21 September 1990*: p. 153-160
- Kuwata, H., Saitoh, H. and Shimada, K. (1994) Occurrence of *Fusarium* wilt of tomato, caused by *Fusarium oxysporum* fsp. *lycopersici* race J2, in Aomori Prefecture, and its control by grafting to rootstocks. *Bulletin of the Aomori agricultural experiment station* 34: p. 1 -10
- Lodha, S. (1995) Soil solarisation, summer irrigation and amendments for the control of *Fusarium oxysporum* fsp. *cumini* and *Macrophomina phaseolina* in soils. *Crop protection* 14(3): 215-219
- Lu, MA Chen, Y.H., Lin, M.S. and Chen, H.P. (1992) An experiment using grafting for control of tomato bacterial wilt. *Plant protection* 18(3): p. 25
- Marois, J.J., Johnston, S.A., Dunn, M.T. and Papavizas, G.C. (1982) Biological control of *Verticillium* wilt: of eggplant in the field. *Plant disease* 66: p. 1166-1168
- Morra, L., Mennella, G. and D'Amore, R. (1992) Innesto della melanzana (*Solanum melongena* L.) quale mezzo per la difesa da patogeni tellurici e l'aumento dell' resa. 11. Contributo. *Culture Protette* 21(12): p. 85-93
- Osmelak, J.A. (1990) Status and management of major vegetable pests in Australia. In: Lim, G.S. and Di, Y.-B. (eds) *Status and management of major vegetable pests in the Asia-Pacific region*. RAPA publication 1990/3: p. 12-22
- Porter, I.J. and Merriman, P.R. (1985) Evaluation of soil solarisation for control of clubroot of crucifers and white rot of onions in Southeastern Australia. In: Parker, C.A., Rovira, A.D., Moore, K.J. and Wong, P.T.W. (eds) *Ecology and management of soil-borne plant pathogens*. American Phytopathological Society St Paul: p. 282-284
- Prasad, D. and Chawla, M.L. (1991) Importance of environmental factors in management of phyto-nematodes. In: Mohan, 1. (ed) *The fragile environment*. New World Environment Series, New Delhi, 5: p. 115-125
- Prasad, J.S. and Reddy, K.V.S. (1994) Hot water treatment for banana planting material made easier. *Informusa* 3(2): p. 16
- Purwati, E. and Hanudin (1995) Screening of some tomato genotypes for resistance to bacterial wilt by IBWDN method in Indonesia. *Proceedings of the AVNET 11 midterm workshop, Los Banos, the Philippines*: p. 127-129
- Qiu, W. F. (1990) Part 1: Status and management of major vegetable pests in the peoples republic of China. In: Lim, G.S. and Di, Y.-B. (eds) *Status and management of major vegetable pests in the Asia-Pacific region*. RAPA publication 1990/3: p. 30-35

- Rangaswamy, S.D. and Reddy, P.P. (1993) Effect of leaf extracts of trap crops on the growth of tomato and development of *Meloidogyna incognita*. Current nematology 4(1): p. 7-10
- Rao, G.N. (1976)
Control of nematodes in nursery soils. Plant. Chron. 71: p. 257, 259
- Sariah, M. and Tanaka, K. (1994)
Effect of flooding on the chili crop in 28th March, Malaysia. In: 4th Malaysia: p. 90-91
- Sethi, C.L. and Gaur, H.S. (1986)
Nematode management and priorities. In: Indian Agricultural Research, New Delhi: p. 424-445
- Takeuchi, S. (1990)
Status and management of major stable pests in Japan of major stable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 46-56
- UNDID (1995)
UNDP survey on major stable pests in the Pacific Region. In: UNDP Report No. 29-2000 Asia 1995): pp. 39.
- Vattanatangum, A. (1990)
Status and management of major stable pests in Thailand of major stable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 91-95
- Vos, J.G.M. and Sumami, N. (1997)
Integrated pest management of fruit pepper (*Capsicum frutescens*) under tropical conditions. In: 72(2): In press
- Vos, J.G.M., Nurtika, N. and Sumami, N. (1993)
A survey of major stable pests in the Philippines. In: 10(2): p. 91-109.
- Vu, V.M. (1990)
Status and management of major stable pests in Vietnam of major stable pests in the Asia-Pacific region. RAPA publication 1990/3: p. 96-100